




Harrisia cactus

Moonlight cactus

Eriocereus martinii and *Eriocereus tortuosus*

DECLARED CLASS 2



Harrisia Cactus is a native of South America and was introduced to Queensland as a pot plant in the 1890s.

Dense infestations choke out pasture. The sharp spines, even in light infestations, make pasture unfavourable to stock and interfere with operations such as mustering.

The plant fruits prolifically and seeds are spread widely by birds and animals. Harrisia cactus can also reproduce by stem sections taking root. A deep, underground, tuberous root system allows the plant to survive catastrophes which may kill the above ground parts.

Description

Harrisia cactus is a perennial. The spiny fleshy stems are jointed and form tangled mats about half a metre high. Many branches often lay flat and take root where they touch the ground. Each section is ribbed lengthwise with six ribs; each rib has low, thick, triangular humps at regular intervals. These humps have cushions of grey felty hairs, three to five short spines lying flat, and one to three erect, stiff, very sharp spines 2.5–3 cm long.

The large flowers open at night. Flowers are pink, funnel shaped with a tinge of white. These grow singly near the ends of the stems on a scaly but spineless slender grey/green tube 12–15 cm long.

Round red fruits 4–5 cm across have scattered bumps with hairs and spines. Numerous small black seeds are embedded in the white, juicy pulp of the fruit which splits open when ripe.

Harrisia cactus roots are of two types. Shallow feeding roots up to 3 cm thick and 30 cm–2 m long grow mostly horizontally off a crown, up to 15 cm below ground level. Swollen tuberous storage roots descend to a depth of 15–60 cm.

The problem

Harrisia cactus can form dense infestations that will reduce pastures to a level unsuitable for stock. Harrisia cactus will choke out other pasture species when left unchecked.

The spines are a problem for stock management, interfering with mustering and stock movement.

Harrisia cactus produces large quantities of seed that is highly viable and easily spread by birds and other animals. As well as reproducing from seed, Harrisia Cactus has long trailing branches that bend and take root wherever they touch the ground. Any broken off portions of the plant will take root and grow.

Control of this plant is difficult as Harrisia Cactus has a deep underground tuberous root system.

Life cycle

Harrisia cactus bears a bright red fruit containing from 400–1000 small black seeds. Fruit and seed are readily eaten by birds and to a lesser extent by feral pigs. Plants are easily established from seed dropped by these animals. Seeds germinate soon after rain.

Seedlings quickly produce a swollen tuberous food storage root that develops as the plant grows. Branches take root where they touch the ground and new plants will grow from broken branches and sections of underground tubers.

Counts of tubers in dense cactus infestations have shown over 125 000 per hectare. Each plant houses many dormant underground buds that are all capable of resprouting when the tip growth dies, any small portion of the tuberous root is left in the soil will grow.


Habitat and distribution

Harrisia cactus is a native of Argentina and Paraguay, South America. It was introduced to Australia as a pot plant in the 1890s. In 1935 it was first recognised as a serious pest in the Collinsville district and by the 1950s was rapidly spreading south.

Harrisia cactus is mainly a pest of brigalow and associated softwood country. However, infestations are now appearing in box and ironbark stands and also in pine forests. The cactus is shade tolerant and reaches its maximum development in the shade and shelter of brigalow scrub, though established infestations can persist once scrub is pulled.

Harrisia cactus is found in the Collinsville, Nebo, Moranbah, Dingo, Blackwater and Goondiwindi districts with minor infestations occurring at Millmerran, Greenmount, Gatton, Ipswich, Rockhampton, Rannes, Mt Morgan, Alpha and Mitchell (see map).

FIGURE 1 – DISTRIBUTION OF HARRISIA CACTUS IN QUEENSLAND



Declaration details

Harrisia cactus is a declared plant under *Land Protection (Pest and Stock Route Management) Act 2002*. Declaration requires landholders to control declared pests on the land and waters under their control. A Local Government may serve a notice upon a landholder requiring control of declared pests.

Control

Mechanical control

Dig out plants completely and burn. Ensure that all tubers that can grow are removed and destroyed.

Ploughing is not considered an effective means of control unless followed by annual cropping.

Biological control

Two introduced insects have become established in the field:

- a stem boring longicorn beetle *Alcidion cereicola*; and
- a mealy bug *Hypogeococcus festerianus*.

The stem-boring beetle only attacks older woody stems. In the Collinsville area large beetle colonies developed and contributed to the collapse of dense areas of cactus. Populations of *Alcidion cereicola* have declined with the reduction in the cactus in recent years.

The most successful biological control agent is the mealy bug *Hypogeococcus festerianus* which is now present in harrisia cactus in Collinsville, Dingo, Moranbah, Blackwater, Nebo, Charters Towers and Goondiwindi districts with small colonies established at Alpha, Capella, Rannes, Gatton, Greenmount, Millmerran and Rockhampton.

How mealy bug works

The mealy bug aggregates and feeds in the tips of stems and buds, where it limits growth and causes distortion. This results in the knotting of the stem. The plants response is to utilise energy reserves within the tuber system to produce new growth. Eventually the plant dies, as it is unable to support the high continuous energy demands.

Dry weather reduces the effectiveness of the mealy bug. When dry the plants tuber system becomes dormant. Consequently, mealy bug damage does not result in new growth and the energy reserves within the plant are not affected. Instead the bug may damage all vegetative parts and eventually die out. The tuber will remain dormant until adequate moisture returns when it will reshoot.

How to spread the bug

Mealy bug disperses naturally via wind although landholder assistance is necessary for its continuous spread, particularly between patches. The bug is manually spread by cutting infected stems and placing them into healthy plants. The best pieces for starting new colonies are large knobs of twisted and distorted cactus that contain many mealy bugs well protected inside knots. Stem tips covered by white, woolly masses of bug are also good. To collect the bug cut infected stems approximately 15 cm from the distorted knob and place segments in green, plump sections of the healthy plant. Avoid placing mealy bug in stressed or dried out stems. Small cactus plants require at least one large knot, with larger plants requiring three knots per plant. Where possible landholders should infest every cactus clump as this ensures a rapid reduction in growth and fruiting potential. When

cactus infestations are light, chemical control may be a preferable option.

Cut pieces can be transported in boxes or open vehicles. They are not delicate but are better kept in the shade. Avoid keeping them in large heaps, direct sunlight, under tarpaulins or in closed containers for long periods. Such conditions will promote rotting of the stems, leading to poor results or failures. Ideally, stems should be put out within 3 days and a maximum of 5 days.

When to infest

Best results come by infesting new areas during spring and early summer, from September to December. Maximum growth and spreading occurs in the summer months of December to February. During the drier and colder months of April to August the mealy bug does not die, but little growth and multiplication occurs. Introduction of mealy bug during autumn and winter will not be lost but little effect is seen until the following summer.

How soon to expect results

As mealy bugs are generally more active and effective on *Harrisia Cactus* growing underneath shrubs and trees, results will be seen more quickly in these areas than in cactus growing in the open. Best results are obtained when infesting plants that have actively growing new shoots.

During wet summers in northern and central Queensland, the growing points of stems will begin to curl after about six weeks.

By the end of the first summer, damage (severe twisting) will be widespread in infested plants. If the initial infestation was sufficiently heavy, no fruit or growth will occur during the second year, and the cactus will begin to die during the third year. Seedlings and regrowth shoots will continue to be present but by the end of the fourth year there should be very little cactus left.

In the southern portion of the State, where temperatures are lower, the mealy bug still provides control but the process takes longer. Also the mealy bug does better on cactus in the open rather than in the shade as temperatures are higher in the open.

Where to obtain mealy bugs

If you cannot obtain mealy bugs from your own property or neighbour contact the vegetation management/weed control/environmental staff at your local government.

Chemical control



Before using any herbicide always read the label carefully. All herbicides must be applied strictly in accordance with the directions on the label.

Further information

Further information is available from the vegetation management/weed control/environmental staff at your local government.

TABLE 1 – HERBICIDES REGISTERED FOR THE CONTROL OF HARRISIA CACTUS

Situation	Herbicide	Rate	Comments
pastures; non agricultural land	Metsulfuron (eg. Aim WDG, Brush-off)	20 g/100 L water + surfactant	Spray plant when actively growing to run-off point A follow-up treatment may be necessary
pastures; non agricultural land	Access	1 L/60 L diesel	Spray plant when actively growing. Apply as overall spray, wetting all areas of the plant to ground level
pastures; non agricultural land	Tordon TCH Tordon DSH	5 L/100 L water 2.5 L/100 L water	Spray plant when actively growing. Treat all stems thoroughly



Fact sheets are available from NRW service centres and the NRW Information Centre phone (07 3237 1435). Check our web site <www.nrw.qld.gov.au> to ensure you have the latest version of this fact sheet. The control methods referred to in this Pest Fact should be used in accordance with the restrictions (federal and state legislation and local government laws) directly or indirectly related to each control method. These restrictions may prevent the utilisation of one or more of the methods referred to, depending on individual circumstances. While every care is taken to ensure the accuracy of this information, the Department of Natural Resources and Water does not invite reliance upon it, nor accept responsibility for any loss or damage caused by actions based on it.